

A WOMAN'S RIGHT TO KNOW

Informational material

Draft for review purposes only.



All rights reserved, 2016. Texas Department of State Health Services. Revised May 2016.
Color pictures representing the development of the child at two-week gestational increments:
Peg Gerrity, www.peggerrity.com.

Introduction

You are pregnant and want to know everything you can about the options you have. You have a right to know the truth. This booklet provides important information about the baby that is growing in your womb and the resources available to you during and after your pregnancy. You need good information in order to make important decisions about your pregnancy and your life. You have the right to make these decisions freely. No one else should make them for you.

If you are reading this because you are already considering an abortion, the doctor who agrees to perform the abortion must wait at least 24 hours before performing the abortion so that you can consider all the facts and make this important decision freely. Only you have the right to decide what to do.

You and your doctor should talk openly and privately. You have the right to ask questions and know as much as you can. The best way for you to understand these risks is to share your health history with your doctor and discuss the risk of each option in light of your personal health history and needs. Some things you should discuss with your doctor include:

- Your personal health history.
- How long you have been pregnant.
- The medical risks of having an abortion.
- The alternatives to abortion, including adoption.
- The medical risks of carrying a pregnancy to term.
- The many public and private agencies that will help new mothers through pregnancy,

childbirth and motherhood, including nearby adoption agencies and free sonogram services, which provide pictures of the baby in your womb.

- Money and other types of support available to new mothers.

Take whatever time you need to read this booklet and talk to other people you trust. You might speak with a family member, a spiritual or professional counselor, a close friend, your spouse, your partner or the father of the baby. You deserve the advice and support of those you trust, and you are strongly urged to ask for their advice and guidance before you make decisions that affect your pregnancy. You will need the support of those closest to you, whatever you decide. The decisions you make about your pregnancy are very important — you have the right to make them based upon your values, your beliefs and your health care needs.

You can view additional materials online at www.dshs.state.tx.us/wrtk.^{*} This website is secure. No one from the Texas Department of State Health Services (DSHS) will collect or record any information about you.

If you are under 18 years old, Texas law requires a doctor to notify your parent or guardian before you can have an abortion. In most cases, the parent or guardian must give consent unless a waiver is given. If you are a minor, ask the doctor or clinic for the booklet, *So You're Pregnant, Now What?*, which discusses this part of the law. This booklet is available at www.dshs.state.tx.us/adolescent/resources.shtm.

**Note: A Woman's Right to Know: Resource Directory (www.dshs.state.tx.us/wrtk) lists information on certain programs and services that can be provided to women during pregnancy, childbirth and as the child is growing up. The booklet contains names, addresses and telephone numbers of these programs. The Resource Directory also has information about public and private adoption agencies.*

“Take the time you need to talk with people that you trust...”

Your baby's development

Fetal Pain

Newborn babies are able to feel pain. We know that babies develop the ability to feel pain while in the womb. In consideration of the potential for fetal pain, Texas law currently limits abortion to under 20 weeks.¹

The next section will follow the changes that will be seen as a baby is developing throughout pregnancy. The period of time when the baby develops inside the mother's body is called gestation.


The development of the baby can be measured in two ways:

- The number of weeks since the start of the last normal menstrual cycle (weeks of gestation).
- The number of weeks since the estimated date of conception, around two weeks after the start of the last normal menstrual cycle.

Usually, your baby's due date is estimated to be 40 weeks after the start of your last normal menstrual cycle.

Week 2 (4 weeks of gestation)

- Your baby's brain and the spinal cord begin to form.
- The heart begins to form.
- The stomach and intestines are forming.
- The bone tissue is growing.
- The baby is scientifically referred to as an embryo.
- Your baby's weight is less than 1 ounce and length is less than $\frac{1}{8}$ inch.



Week 4 (6 weeks of gestation)

- Your baby's developing heart starts to beat.
- The lungs, liver, stomach and other major organs begin to form.
- Brain activity can be recorded, and the brain and spinal cord are completing development.
- The arm and leg buds (the structures that will become the limbs) are present.
- The eyes are present.
- Your baby is less than $\frac{1}{4}$ inch long.



Week 6 (8 weeks of gestation)

- Your baby's developing heart beats with a regular rhythm.
- Facial features — the eyes, nose, lips and tongue — start to develop.
- All essential organs have begun to form.
- The spinal nerves begin to develop.
- The brain continues to form and begins to control organs.
- The arms and legs grow longer.
- Fingers and toes are present.
- Sex organs are beginning to form.
- Your baby is 1/2 inch long.



Week 8 (10 weeks of gestation)

- Your baby has his or her first spontaneous movements (movements that happen on their own).
- Elbows are formed and fingernails appear.
- The eyelids are more developed.
- The external ears begin to take final shape.
- Facial features continue to develop.
- After the end of this week, the baby is scientifically referred to as a fetus.
- Your baby is about 1 ¼ to 1 ½ inches from head to bottom.



Week 10 (12 weeks of gestation)

- Your baby moves, but you cannot feel the movements.
- The fibers that carry pain to the brain are developed.
- All the body parts and organs are formed.
- The heartbeat can be heard with electronic devices.
- The body grows rapidly.
- The body grows longer, and the neck lengthens.
- The eyelids close and will not reopen until about the 28th week.
- Fingernails appear.
- Teeth buds begin to form.
- Your baby weighs about half an ounce and is about 2 1/2 inches long from head to bottom.



Your baby's development

Week 12 *(14 weeks of gestation)*

- Your baby's mouth makes sucking motions.
- The taste buds are developing.
- The arms and legs begin to move.
- Hiccup movements are present.
- The external sex organs are developed and are clearly either male or female.
- Your baby is about 3½ inches long from head to bottom and weighs about 1½ ounces.



Week 14 *(16 weeks of gestation)*

- Your baby's swallowing and chest movements are clearly present.
- Hand-to-face movements are common.
- The eyes are beginning to move.
- Your baby is about 4¾ inches long from head to bottom and weighs less than 4 ounces.



Week 16 *(18 weeks of gestation)*

- Your baby's arms and legs begin to punch and kick, and you may feel these movements.
- The kidneys are functioning and make urine.
- Taste buds are present.
- The skin is wrinkled.
- Features further develop — eyelids, ear and upper lip.
- Your baby is about 5½ inches long from head to bottom and weighs about 7 ounces.



Week 18 (20 weeks of gestation)

- Your baby is more active, and you can feel the movement, including turning side to side and front to back.
- Breathing-like movements become regular and are detected by ultrasound.
- The structures of the ears are well-developed.
- Ovaries containing eggs have formed in females and testes begin to descend in males.
- Meconium (a greenish mixture of swallowed cells and secretions) begins to form in the intestinal tract. This will be the baby's first bowel movement.
- Your baby weighs about 11 ounces and is about 6 inches from head to bottom.



Week 20 (22 weeks of gestation)

- Your baby is beginning to develop regular sleeping and waking patterns.
- The eyelids and eyebrows are well formed, and eyes are fully functional.
- The vocal cords are active, and reflexes are present.
- The heartbeat can be heard with a stethoscope.
- The body is covered with fine hair called lanugo.
- Your baby weighs about 1 pound and is about 7 to 7 ½ inches in length from head to bottom.



Week 22 (24 weeks of gestation)

- Your baby will blink and startle (pull in arms and legs) if stimulated.
- Skin is wrinkled and red.
- The senses of smell and taste are developed.
- The inner ear has reached adult size, and the baby may hear some sounds.
- A baby born at this time will attempt to breathe, but the lungs are not fully developed.
- Your baby weighs about 1.4 pounds and is about 8 inches from head to bottom.



Your baby's development

Week 24 (26 weeks of gestation)

- Your baby's lungs are now fully formed but are not yet ready to function outside the uterus.
- The lines on the skin of the fingers (fingerprints), toes, palms of the hands and soles of the feet are now formed.
- Your baby weighs about 1.8 pounds and is about 9 inches from head to bottom.



Week 26 (28 weeks of gestation)

- Your baby's brain is fully formed, and the nervous system continues to develop.
- The lungs and digestive system are fully formed and continue to develop.
- The eyes are partially open and eyelashes are present.
- Your baby weighs about 2.2 pounds and is almost 10 inches from head to bottom.



Week 28 (30 weeks of gestation)

- Your baby kicks and stretches.
- Your baby makes grasping motions and responds to sound.
- The nervous system controls some body functions.
- Your baby weighs about 2.9 pounds and is more than 10 ½ inches from head to bottom.



Week 30 (32 weeks of gestation)

- Your baby gains weight, and the body fills out.
- Toenails and fingernails start to grow.
- The pupils will react to light and constrict (become smaller).
- Your baby weighs about 4 pounds and is almost 11 inches from head to bottom.



Week 32 (34 weeks of gestation)

- Your baby is active (moving) 60 percent or more of the time.
- Bones harden, but the skull remains soft and flexible for delivery.
- There are rhythmic breathing movements, but the lungs are still developing.
- Your baby weighs about 4.6 pounds and is almost 12 inches from head to bottom.



Week 34 (36 weeks of gestation)

- Your baby rapidly gains weight in preparation for birth.
- The body and face fill out; the skin no longer appears wrinkled.
- The fingernails reach the ends of the fingertips.
- Large amounts of immune factors (substances in the blood that protect against disease or infection) are passed from you to your baby during the last four gestational weeks.
- Your baby weighs about 5.5 pounds and is about 12 ½ inches from head to bottom.



Week 36 (38 weeks of gestation)

- Your baby's lungs and brain are developing rapidly.
- Your baby can grasp firmly.
- Your baby weighs about 6.4 pounds and is about 13 1/3 inches from head to bottom.



Week 38 (40 weeks of gestation)

- Your baby is fully developed and considered full-term.
- The brain and nervous system will continue to develop after birth.
- The toenails have reached the tips of the toes.
- Small breast buds are present.
- The fingernails extend beyond fingertips.
- Your baby weighs about 7.5 pounds and is about 14 inches from head to bottom.



Abortion risks

This section describes the risks associated with an abortion. The risks of having an abortion can vary depending on several factors.

Death

You have a greater risk of dying from the abortion procedure and having serious complications the further along you are in your pregnancy. There is one death per every one million abortions for women who are eight weeks pregnant or less, one death per 29,000 abortions for pregnancies at 16 to 20 weeks of gestation, and one death per 11,000 abortions at 21 weeks of gestation and later.ⁱⁱ

Mental Health Risks from Abortion

Women report a range of emotions after an abortion. This can include depression or thoughts of suicide. Some women have also reported after their abortion feelings of grief, anxiety, lowered self-esteem, regret, sexual dysfunction, avoidance of emotional attachment, flashbacks and substance abuse. For some women, these emotions may appear immediately after an abortion or gradually over a longer period of time. These feelings may recur or be felt more strongly at the time of another abortion, a normal birth or on the anniversary of the abortion. It is important that you talk to your doctor if you experience these feelings.

Counseling or support before you make a decision to have an abortion is very important. If family help and support are not available to you, talking with a spiritual or professional counselor before having an abortion can help you better understand your decision. Many pregnancy resource centers can provide counsel to you. These centers are listed in the resource directory.



Future Infertility

The further along you are in your pregnancy, the greater the chance of serious complications that can cause you to be infertile and the greater the risk of dying from the abortion procedure. Some complications associated with an abortion, such as an infection, a cut or a torn cervix, may make it difficult or impossible to become pregnant in the future or to carry a pregnancy to term.

Breast Cancer Risk

Your pregnancy history affects your chances of getting breast cancer. If you give birth to your baby, you are less likely to develop breast cancer in the future. Research indicates that having an abortion will not provide you this increased protection against breast cancer. In addition, doctors and scientists are actively studying the complex biology of breast cancer to understand whether abortion may affect

the risk of breast cancer. If you have a family history of breast cancer or breast disease, ask your doctor how your pregnancy will affect your risk of breast cancer.

Other Physical Risks

Additionally, abortion could result in physical side effects, with different levels of severity. A woman will usually have cramping and vaginal bleeding after any type of abortion procedure. Other symptoms or side effects include nausea (feeling sick to your stomach) or vomiting, diarrhea, warmth or chills, headache, dizziness and fatigue (feeling very tired).

Complications of spontaneous miscarriages and elective abortions can result in injuries to the internal organs, blood clots or serious infections. These will be listed in detail for each type of abortion later in this booklet.



Making an informed decision

You need to know as much as you can about your options in order to make an informed decision. Each option has possible risks and benefits. There are counseling services available to help you fully understand your options and make your decision.

Before an Abortion

You should ask your doctor about any risks you might face. It is your right and your doctor's responsibility to make sure that you are fully informed before deciding to have an abortion. Here are some things the doctor should talk to you about:

- You will find out for sure if you are pregnant and how long you have been pregnant. Your doctor will also do a pelvic exam.
- Your doctor will evaluate your health and discuss medical risks of having an abortion. You will have a physical exam, be asked about your medical history and have lab tests done. The doctor will answer any questions you might have.
- Your doctor will describe the baby growing in your womb to you and give you a list of agencies that offer alternatives to abortion, including adoption, and the names of agencies and private counseling organizations that provide ultrasound services (images of the baby in the womb or sonogram). Some organizations provide ultrasounds for free.
- If you decide to have an abortion, at least 24 hours before an abortion is performed, the doctor will give you an ultrasound and tell you what you are seeing, including the size of your baby and any heartbeat, limbs and internal organs. The doctor is required to show you the ultrasound picture while it is being performed. You may choose not to view these images of your baby. You have the right to view your ultrasound at any time.

- If you are under the age of 18, a parent or guardian must be notified and agree to the abortion, or you will have to ask a judge to waive that notification and consent requirement.
- The doctor performing the abortion must provide you with the Texas Department of State Health Services (DSHS) *A Woman's Right to Know* printed materials and let you know that the materials are available on the DSHS website. You will have at least a full day to read the information your doctor gives you before the appointment for your abortion. If you decide to have the abortion, you must sign a written consent for that abortion.
- Your doctor must also provide you with a telephone number that you can call 24 hours a day to talk with the doctor or other healthcare personnel about any complications or questions related to the abortion and the name and telephone number of the hospital that is closest to your home where you could be treated in the case of an abortion-related emergency.
- Your doctor may offer you a medical abortion, which is a medication given to cause an abortion. If a medical abortion is performed, your doctor must provide you with a copy of the final printed label of any abortion-inducing drugs used in the procedure.



Medical and Social Assistance

Your doctor must give you certain important information before the abortion can be done, including:

- Medical assistance benefits that can help with prenatal care, childbirth and neonatal care.
- The legal responsibilities of the baby's father in helping to support your child if you decide to stay pregnant and keep the baby. The law says he must help even if he offered to pay for an abortion.
- Public and private agencies can help you if you were raped or a victim of incest.
- Public and private agencies can give you information about preventing pregnancies and medical referrals for birth control methods.

You should know that if you choose to have your baby and find yourself weighed down by the job of being a parent, Texas has the "Baby Moses/Safe Haven" law. The law allows you or the baby's father to leave a baby who appears to be under 60 days old in the care of a designated emergency care provider such as an emergency medical services station, fire

station, licensed child-placing agency or any hospital. You do not have to return for the baby, and you will not be charged with a crime if the baby is unharmed.

Child Support Services

Assistance in Obtaining Child Support

Your child's father is legally required to pay to support the child, and the Texas Office of the Attorney General can assist you in obtaining this support. The Attorney General's Child Support Division can help locate missing fathers, legally prove who the father is, initiate child support orders, and collect child support payments. Texas is the top-performing state in terms of ensuring parents pay what they owe, with more than \$3.9 billion disbursed to families in 2015 alone. Texas has a high collection success rate: 65.2% of all child support amounts due are collected. If you need services, call 800-252-8014 or visit www.texasattorneygeneral.gov/cs. If you are a survivor of family violence, there are steps you can take to pursue child support safely.

For more information, visit www.getchildsupportsafely.org.





Draft for review purposes only.

***Ask your doctor for a copy of the
“A Woman’s Right to Know: Resource Directory”.***

Adoption Services

Another option to consider is adoption. Adoption means you, as the birth parent are voluntarily transferring your rights as the parent of your baby to another family. Choosing adoption means you want your baby to have a good life, but right now may not be the best time for you to be a parent.

Adoption is a brave, loving choice for your baby. When you place your baby for adoption, you are placing him or her with loving parents who can raise your baby to have the good life that you want for your baby.

There are many resources available to help you decide whether adoption is the right choice for you and your baby. Each adoption is different, and help is available to make sure that the adoption process fits you and your baby's needs.

If you choose adoption, you may be able to select and meet the family who will adopt your

baby. There are two different types of adoptions that you can choose for your baby. An open adoption allows you to stay in contact with your baby as he or she grows up, perhaps through pictures, phone calls or visits. A closed adoption means there will be no contact between you and the family adopting your baby once the adoption is finalized. What type of adoption you want is entirely up to you.

Talk with a family member, a spiritual or professional counselor, or a close friend to seek their advice on adoption. Since adoption is a big decision, you may want the support of those you trust when making this decision.

It is never too late to choose adoption for your baby. You can make this choice anytime during your pregnancy or even after your baby has been born.

You can find more information about adoption and organizations that offer adoption services at: <https://www.dshs.state.tx.us/wrtk/resources>.



Abortion procedures and side effects

The types of abortion procedures include medical abortions and surgical abortions. A woman will usually have cramping and vaginal bleeding after any type of abortion procedure. There are other symptoms or side effects that may occur with any type of abortion procedure. These include nausea (feeling sick to your stomach) or vomiting, diarrhea, warmth or chills, headache, dizziness and fatigue (feeling very tired). Also, other serious complications sometimes arise. This section will also discuss the risk of complications for each of the methods.

First Trimester Abortions

During the first trimester (through 13 weeks of gestation) an abortion can be performed through medicine or surgery.

Medical (Nonsurgical) Abortion

Medical abortion uses medicine to end a pregnancy instead of surgery and is used early in pregnancy — up to 70 days (10 weeks) from the first day of your last menstrual period. This method requires several visits to your doctor. The medicines used for a medical abortion cause bleeding, cramping and passing of the fetus and other tissue. In some cases, excessive bleeding may require blood transfusions, treatment with medication, surgery or saline transfusions. Severe infection is a known risk following a medical abortion.

Possible complications or risks of a medical abortion using mifepristone and misoprostol pills:

- About 5 to 8 percent of procedures will not work and will need a surgical procedure to end the pregnancy.ⁱⁱⁱ
- Hemorrhaging (heavy bleeding) with possible emergency hysterectomy (removal of the uterus) to end the bleeding.
- About 1 of every 100 medical abortions will require a surgical procedure of some kind to stop severe bleeding.ⁱⁱⁱ

- Failure to remove all parts of the baby and other tissue, including the placenta, which may require another procedure.
- Increased risk of infertility (the inability to have a baby), if complications occur with the procedure.
- Infection, which is usually caused by an infection the woman already had at the time of the abortion. Serious infection can also occur and has resulted in death in a small number of cases.



Who should not have a medical abortion?

Some women should not have a medical abortion. Some reasons a medical abortion would not be done include:

- It has been more than 70 days (10 weeks) since the first day of the last menstrual period.
- The woman is allergic to one of the medications to be used.
- The woman has or might have a tubal or ectopic pregnancy (where the baby grows outside of the uterus).
- The woman has an intrauterine contraceptive device (IUD). The IUD must be taken out before a doctor can give a woman the medicine.
- The woman is taking certain types of medicines, like blood thinners or certain steroids.
- The woman has a particular medical condition, like uncontrolled seizures or a bleeding disorder.
- The woman cannot get to all of the visits needed to get the medicines to complete the abortion or cannot get to the checkup(s) needed after the abortion.
- The woman cannot get to emergency care if needed in the two weeks after taking the medicines.

- The woman has problems with her adrenal glands (chronic adrenal failure).

You should discuss with your doctor whether you have any medical condition or other problems that would make a medical abortion unsafe for you.

When having a medical abortion, your doctor will give you mifepristone (also called RU 486) during an office visit. It is taken by mouth. After receiving mifepristone, you will have vaginal bleeding and pass clots and fetal tissue. Bleeding usually lasts 9 to 16 days and may last up to 30 days. Your doctor will make an appointment for two days after you take mifepristone. If the medical abortion is not complete, you will be given a second drug, misoprostol. Misoprostol may cause cramps, nausea, diarrhea and other symptoms. Your doctor may send you home with medicines for these symptoms.

Your doctor will make a follow-up appointment for you within 14 days of the day you take the medication. Your doctor will check your health for any symptoms you may be experiencing, as well as whether your pregnancy has completely ended. Make sure to discuss any physical and emotional symptoms you are experiencing with your doctor.



Abortion procedures and side effects

First Trimester Abortions (cont.)

Suction Curettage (Surgical Abortion)

The most common type of abortion is the suction curettage, sometimes called a dilation and curettage, or vacuum curettage. After the first trimester, this procedure is called dilation and evacuation.

Most women will have some pain with this procedure. Before the procedure, you may be given pain medication, a sedative or both. For the procedure, you may receive local anesthesia injected or applied in the area of the cervix. You may also receive general anesthesia that will put you to sleep so that you do not feel pain during the procedure.

The doctor inserts a speculum to hold open the vagina, as is done for a pelvic exam, and injects or applies local anesthesia. The cervix is stretched open. Then the contents of the uterus, including the baby and placenta, are removed using a suction device that is inserted into the uterus.

The procedure usually takes 10 to 15 minutes, but can take longer depending upon the growth stage of the baby.

Possible complications or risks of suction curettage include:

- Hemorrhaging (heavy bleeding) with possible emergency hysterectomy (removal of the uterus) to end the bleeding
- Perforation of the uterus (a hole in the uterus)
- Injury to the bowel or bladder, if there is a perforation of the uterus
- Abdominal incision and operation to correct injury
- Failure to remove all parts of the baby and

other tissue, including the placenta, which may require another procedure (occurs in fewer than 50 per 1,000 procedures, or less than 5 percent^{iv})

- Increased risk of infertility (the inability to have a baby), if complications occur with the procedure
- Infection, which is usually caused by an infection the woman already had at the time of the abortion



Second Trimester Abortions

During the second trimester (between 13 weeks of gestation and 22 weeks of gestation) an abortion can be performed by a surgical method known as dilation and evacuation.*

Dilation and Evacuation (D&E)

Most second trimester abortions are done by dilation and evacuation (D&E). This surgical procedure is usually performed in a surgical center or hospital.

Before this procedure, the doctor will prepare the cervix. This process softens and stretches open the cervix and may require one or more visits.

The softening and opening of the cervix before performing the D&E helps decrease the risk of tears or lacerations to the cervix.

Most women will experience some pain with this procedure. Before the procedure you may be given pain medicine, a sedative or both. For the procedure, you may receive local or regional anesthesia injected or applied in the area of the cervix. You may also receive general anesthesia that will put you to sleep during the procedure. Your doctor will discuss your options, and any risks, for anesthesia.

At the beginning of the procedure, the doctor will make sure that the cervix is open. The fluid surrounding the baby (amniotic fluid) will be removed with a suction device placed into the uterus. The baby and placenta are removed from the uterus using surgical instruments. Finally, a suction device will be inserted into the uterus at the end of the procedure to remove

any fetal tissue that remains. After 14 weeks of pregnancy, the baby, placenta and other contents of the womb may be removed in pieces using surgical instruments. This procedure usually takes less than one hour.

Possible complications or risks of a D&E include:

- Death, in rare cases
- Cervical laceration (tears or injury to the cervix)
- Hemorrhaging (heavy bleeding) with possible emergency hysterectomy (removal of the uterus) to end the bleeding
- Perforation of the uterus (a hole in the uterus)
- Injury to the bowel or bladder, if there is a perforation of the uterus
- Abdominal incision and operation to correct injury
- Failure to remove all parts of the baby and other tissue, including the placenta, which may require another procedure
- Increased risk of infertility (the inability to have a baby), if complications occur with the procedure
- Embolism (a loose blood clot or amniotic fluid, fetal cells, fetal hair or other matter in the bloodstream that causes sudden blockage of blood flow to the lungs or other organ)

** Note: Texas law states that abortion may not be performed after the post-fertilization age of 20 weeks or more, which is approximately 22 weeks or more of gestation, as gestation is defined in this booklet. Texas law allows for exceptions when the life of the mother is threatened, serious risk exists of irreversible impairment of a major bodily function (other than a psychological condition) or a severe fetal abnormality is present.*

Pregnancy and Childbirth

Birth is a life-changing experience, and each birth brings a new and different set of experiences and feelings. Pregnancy and birth is usually a safe and natural process, although complications can occur.

There are complications associated with pregnancy and childbirth. The most common complications of pregnancy include:

- Tubal or ectopic pregnancy (where the baby grows outside of the uterus)
- High blood pressure
- Complicated delivery
- Premature labor
- Depression
- Infection
- Diabetes
- Hemorrhage (heavy bleeding)

During labor, the uterus contracts and pushes to deliver the baby. The baby may be delivered through the woman's vagina or by a surgical procedure called cesarean section or c-section.

Vaginal Delivery

Possible side effects and risks

- Injury to the bladder or rectum
- A hole (fistula) between the bladder and vagina or the rectum and vagina
- Hemorrhage (heavy bleeding)
- Infertility (the inability to have a baby) due to infection or complications
- Emergency treatment for any of the above problems, including the possible need to treat with an operation, medicine or blood transfusions
- Death (very rare — one per 500,000 vaginal deliveries of a live-born infant^v)

Cesarean Birth

Possible side effects and risks

- Injury to the bowel or bladder
- Infertility (the inability to have a baby) due to infection or complications
- Hemorrhage (heavy bleeding)
- Injury to the tube (ureter) between the kidney and bladder
- A possible hysterectomy (removal of the uterus) due to complications or injuries
- Complications from anesthesia such as respiratory problems, headaches or drug reactions
- Emergency treatment for any of the above problems, including the possible need to treat with an operation, medicine or blood transfusions
- Death (very rare — one per 45,500 cesarean deliveries of a live-born infant^v)



Postpartum Symptoms

The feelings you experience after birth may be the most intense you have ever encountered: great surges of joy and happiness, feelings of contentment and fulfillment. It is not uncommon for women to also experience fears, worries or sadness. Depression can occur after the birth of a baby, and many new mothers experience various degrees of it. While depression can occur within days after delivery, it can also appear gradually, and sometimes it doesn't start until a year after your baby's birth.

In most cases, mothers have mild symptoms that may last only a few days. However, some mothers experience severe symptoms that can include exhaustion, feelings such as worthlessness or hopelessness, and memory loss. In rare circumstances you may have a fear

of harming yourself or your baby. Although these symptoms may not last long, if you have any of them, you should call your doctor right away and discuss these symptoms. Doctors can give you professional help and support during this time.

You can reduce the risk for problems or complications in any pregnancy by:

- Getting early and regular prenatal care.
- Eating a well balanced diet and get regular exercise.
- Avoiding tobacco, alcohol, or drugs that your doctor hasn't prescribed for you.

If you have questions or concerns, be sure to talk with your doctor or other health care provider.



After an abortion

Call the clinic or doctor that performed the abortion or go to the emergency room if:

- Heavy bleeding occurs (two or more thick pads per hour for two hours in a row) or if you are concerned about heavy bleeding.
- You have stomach pain or discomfort, or feel sick (such as feeling weak or having an upset stomach, nausea, vomiting or diarrhea) with or without fever for more than 24 hours after taking misoprostol for a medical abortion.
- You have a fever (a temperature greater than 100.4 degrees Fahrenheit or 38 degrees Celsius).
- You have difficulty breathing or have shortness of breath.
- You have chest pain.

- Pain is severe or not controlled by pain medication.
- You are disoriented.

Your doctor will schedule one or more follow-up visits, usually beginning within two to three weeks after the procedure. Your doctor must schedule a follow-up visit within 14 days of a medical abortion. It is important that you keep all of your appointments.

You should also consider seeking counseling if you are experiencing depression, thoughts of suicide or other psychological distress following your abortion.



i Bellieni, C. V. (2012). Pain Assessment in Human Fetus and Infants. *The AAPS Journal*, 14(3), 456–461. <http://doi.org/10.12248-012-9354-5>. Last accessed June 6, 2016

ii Bartlett LA, Berg CJ, Shulman HB, Zane SB, Green CA, Whitehead S, Atrash HK. [Maternal and Infant Health Branch, Division of Reproductive Health, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention.] Risk factors for legal induced abortion-related mortality in the United States. *Obstetrics & Gynecology*. 2004;103(4):729-37. <http://www.ncbi.nlm.nih.gov/pubmed/15051566>. Last accessed June 6, 2016

iii U.S. Food and Drug Administration. Mifeprex Medication Guide. 2009. <http://www.fda.gov/downloads/Drugs/Drugsafety/ucm088643.pdf>. Last accessed June 6, 2016.

iv Royal College of Obstetricians and Gynaecologists (RCOG). *The care of women requesting induced abortion*. London (England): Royal College of Obstetricians and Gynaecologists (RCOG); 2011 Nov. 130 p. (Evidence-based Clinical Guideline; no. 7). <http://www.rcog.org.uk/womens-health/clinical-guidance/care-women-requesting-induced-abortion>. Last accessed June 6, 2016.

v Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Pregnancy Mortality Surveillance System. *Pregnancy-Related Mortality in the United States*. Available: <http://www.cdc.gov/reproductivehealth/MaternalInfantHealth/PMSS.html>. Last Accessed 6/20/2016.

Draft for review purposes only.

www.dshs.state.tx.us/wrtk/default.shtm



Texas Department of State Health Services
Office of Title V and Family Health

Publication number 1-450 • Revised May 2016 .

